5

10

15

CLAIMS

What is claimed is:

1. A compound comprising


or a pharmaceutically acceptable salt or a prodrug thereof;

wherein a dashed line represents the presence or absence of a double bond or a triple bond;

A is $-(CH_2)_6$ -, cis $-CH_2CH=CH-(CH_2)_3$ -, or $-CH_2C=C-(CH_2)_3$ -, wherein 1 or 2 carbon atoms may be substituted with S or O;

Z is O, S, or NR;

X is selected from the group consisting of CO₂H, CONHR₂, CONR₂, CON(OR)R, CON(CH₂CH₂OH)₂, CONH(CH₂CH₂OH), CH₂OH, P(O)(OH)₂, CONHSO₂R, SO₂NR₂, SO₂NHR, and




20

25

J is C=O or CHOH;

R is independently H, C_1 - C_6 alkyl, phenyl, or biphenyl; and E is C_3 - C_6 alkyl, C_4 - C_{10} cycloalkyl, phenyl or napthyl having from 0 to 2 substituents, or a heteroaromatic moiety having from 0 to 2 substituents, wherein said substituents comprise up to 4 non-hydrogen atoms.

2. The compound of claim 1 comprising



- 5 or a pharmaceutically acceptable salt or a prodrug thereof.
 - 3. The compound of claim 1 comprising

or a pharmaceutically acceptable salt or a prodrug thereof.

- 4. The compound of claim 3 wherein Z is O.
- 10 5. The compound of claim 3 wherein Z is S.
 - 6. The compound of claim 3 wherein Z is NH.
 - 7. The compound of claim 3 wherein J is C=O.
 - 8. The compound of claim 3 wherein J is CHOH.
 - 9. The compound of claim 3 wherein X is CO_2H .
- 15 10. The compound of claim 3 wherein E is phenyl, thienyl, furyl, pyridinyl, napthyl, benzothienyl, or benzofuryl having from 0 to 2 substituents comprising up to 4 non-hydrogen atoms.
 - 11. The compound of claim 3 wherein E is n-butyl.
 - 12. The compound of claim 1 comprising

20

or a pharmaceutically acceptable salt or a prodrug thereof.

13. The compound of claim 1 comprising

or a pharmaceutically acceptable salt or a prodrug thereof.

14. The compound of claim 1 comprising

5

or a pharmaceutically acceptable salt or a prodrug thereof.

10 15. A liquid composition comprising

or a pharmaceutically acceptable salt or a prodrug thereof;

wherein a dashed line represents the presence or absence of a double bond or a triple bond;

A is $-(CH_2)_6$ -, cis $-CH_2CH=CH-(CH_2)_3$ -, or $-CH_2C\equiv C-(CH_2)_3$ -, wherein 1 or 2 carbon atoms may be substituted with S or O;

Z is O, S, or NR;

X is selected from the group consisting of CO₂H, CONHR₂, CONR₂, CON(OR)R, CON(CH₂CH₂OH)₂, CONH(CH₂CH₂OH), CH₂OH, P(O)(OH)₂,

20 CONHSO₂R, SO₂NR₂, SO₂NHR, and

5

10

15

20

25

J is C=O or CHOH;

R is independently H, C₁-C₆ alkyl, phenyl, or biphenyl; and E is C₃-C₆ alkyl, C₄-C₁₀ cycloalkyl, phenyl or napthyl having from 0 to 2 substituents, or a heteroaromatic moiety having from 0 to 2 substituents, wherein said substituents comprise up to 4 non-hydrogen atoms; wherein said liquid is formulated for ophthalmic use.

16. A method comprising administering a compound to a mammal, wherein said method is useful for the treatment of glaucoma or ocular hypertension in said mammal, said compound comprising

or a pharmaceutically acceptable salt or a prodrug thereof;

wherein a dashed line represents the presence or absence of a double bond or a triple bond;

A is $-(CH_2)_6$ -, cis $-CH_2CH=CH-(CH_2)_3$ -, or $-CH_2C\equiv C-(CH_2)_3$ -, wherein 1 or 2 carbon atoms may be substituted with S or O;

Z is O, S, or NR;

X is selected from the group consisting of CO₂H, CONHR₂, CONR₂, CON(OR)R, CON(CH₂CH₂OH)₂, CONH(CH₂CH₂OH), CH₂OH, P(O)(OH)₂, CONHSO₂R, SO₂NR₂, SO₂NHR, and

J is C=O or CHOH;

R is independently H, C₁-C₆ alkyl, phenyl, or biphenyl; and

E is C₃-C₆ alkyl, C₄-C₁₀ cycloalkyl, phenyl or napthyl having from 0 to 2 substituents, or a heteroaromatic moiety having from 0 to 2 substituents, wherein said substituents comprise up to 4 non-hydrogen atoms.